

On the non-existence of L-space surgery structure

Motoo Tange*

Abstract

We exhibit homology spheres which never yield any lens spaces by integral Dehn surgery by using Ozsváth and Szabó's contact invariant.

1 Introduction

Let Y be a closed oriented 3-manifold. In this paper we write by $Y_r(K)$ the Dehn surgered manifold of a knot K in Y and with slope r . Lens spaces can be obtained from the Dehn surgery of the unknot U with slope $-p/q$, i.e. $L(p, q) = S^3_{-p/q}(U)$.

In general it is difficult to determine when a lens space can be obtained by an integral surgery of a non-trivial knot K in S^3 . There are some well-known non-trivial knots in S^3 yielding lens spaces by integral surgeries, for example torus knots, 2-cable knots of torus knots, and $(-2, 3, 7)$ -pretzel knot and so on.

If we weaken the ambient space to any homology sphere, we can construct many lens spaces by integral Dehn surgery. For example in [1] R. Fintushel and R. Stern have asserted that a lens space $L(p, q)$ is obtained by an integral Dehn surgery of a knot K in a homology 3-sphere Y if and only if $q = \pm x^2 \bmod p$ holds for an integer x . What kinds of homology spheres and the knots can realize for the solution x ? Are such homology spheres and knots restricted? These questions are quite natural ones.

In the present paper we shall show that such homology spheres are restricted (see Theorem 1.1). We denote by $\overline{\Sigma(2, 3, 5)}$ the Poincaré homology sphere with the reversed-orientation as the usual one.

Theorem 1.1 $\overline{\Sigma(2, 3, 5)}$ does not yield any lens spaces by any positive integral Dehn surgeries.

From contact topological view point we would like to consider this theorem.

We explain the original motivation for this paper. In the author's previous paper [9] many lens spaces have been constructed by positive Dehn surgeries on $\Sigma(2, 3, 5)$. But from L-space homology spheres whose correction terms are neither 0 nor 2, the author could not construct any lens spaces by any positive surgeries in some range of the order p of the 1st homology group and correction terms of homology spheres. In particular from $\overline{\Sigma(2, 3, 5)}$ the author could not construct any lens spaces in some range of p . The notions of L-space and correction term shall be defined in Section 2.

¹The author was supported by COE program of Mathematical Department of Osaka University.

On the other hand J.B. Etnyre and K.Honda in [2] have shown that there do not exist any tight contact structures over $\overline{\Sigma(2, 3, 5)}$. Hence I wondered if the two phenomena are related to each other. This is main motivation of this paper.

In [9] the author has studied examples yielding lens spaces by using the surgery formula between invariants of Y , $Y_0(K)$, and $Y_p(K)$. Viewing the examples the author guesses if $Y \in \{\#^n \Sigma(2, 3, 5) \#^m \overline{\Sigma(2, 3, 5)}\}$ carries a positive lens surgery structure, then Y is S^3 or $\Sigma(2, 3, 5)$.

2 Lens space (or L-space) surgery structure

We call a rational homology 3-sphere *L-space* if the Heegaard Floer homology of the manifold is isomorphic to that of S^3 . The definition of Heegaard Floer homology is in [5] and [6]. It is well-known that the set of L-spaces contains all spherical manifolds and some hyperbolic manifolds. In order to generalize Theorem 1.1 to L-space surgery, we need irreducibility of $Y - K$. In fact the trivial 1-surgery of $\overline{\Sigma(2, 3, 5)}$ gives rise to L-space clearly.

The coefficients of the homologies are always $\mathbb{Z}_2$ hence $HF^+(Y, \mathfrak{s})$ is a $\mathbb{Z}_2[U]$ -module, where U is the action to lower the degree of the homology by 2. When Y is a rational homology sphere, the homology admits the absolute $\mathbb{Q}$ -grading as in [4]. The *correction term* $d(Y, \mathfrak{s})$ is defined to be the minimal grading in the $HF^\infty(Y, \mathfrak{s})$ -part, which is the image by the natural map $\pi_* : HF^\infty(Y, \mathfrak{s}) \rightarrow HF^+(Y, \mathfrak{s})$ defined in [5].

Definition 2.1 *Let Y be a closed oriented 3-manifold. We say that Y carries positive (negative) L-surgery structure, if p is a positive (negative) integer and $Y_p(K)$ is an L-space for a null-homologous knot $K \subset Y$. Moreover if $Y - K$ is irreducible, we say that Y admits proper L-surgery structure.*

In particular we say that Y carries positive (negative) lens surgery structure if $Y_p(K)$ is a lens space for a positive (negative) integer p .

We suppose that any connected-sum component of Y is not a lens space. If Y carries positive (or negative) lens surgery structure, then obviously Y carries positive (or negative) proper L-surgery structure.

S^3 carries both positive and negative lens surgery structure. The homology sphere $\Sigma(2, 3, 5)$ carries positive lens surgery structure (see [9]). But for any homology sphere lens surgery structure does not always exist. We shall prove the following in Section 4.

Theorem 2.1 *Let Y be an L-space homology sphere. If Y carries positive proper L-surgery structure, then Y admits positive tight contact structure.*

Let Y be an L-space homology sphere. If $Y_p(K)$ is an L-space, then $\widehat{HFK}(Y, K, g) \cong \mathbb{Z}_2$ holds by using the same method as [8]. It follows from this fact and Y. Ni's result in [11] that if $Y - K$ is irreducible then K is a fibered knot. The tight contact structure above is the one induced from the open book decomposition along the knot K by the method [10] of W. Thurston and H. Winkelnkemper.

From Theorem 2.1 immediately we can prove the following non-existence result for $\overline{\Sigma(2, 3, 5)}$.

Corollary 2.1 *$\overline{\Sigma(2, 3, 5)}$ does not carry positive proper L-surgery structure.*

As we mentioned in Section 1, $\overline{\Sigma(2, 3, 5)}$ carries non-proper positive L-surgery structure.

Proof) Suppose that $\overline{\Sigma(2, 3, 5)}$ carries positive proper L-surgery structure. From Theorem 2.1 $\overline{\Sigma(2, 3, 5)}$ must admit a positive tight contact structure. But $\overline{\Sigma(2, 3, 5)}$ does not admit any positive tight contact structures by the work [2] by J. B. Etnyre and K. Honda. This is inconsistent. $\square$

Proof of Theorem 1.1) It follows from Corollary 2.1 and irreducibility of any lens space that this theorem is true. $\square$

Corollary 2.2 *The homology sphere $\Sigma(2, 3, 5) \# \overline{\Sigma(2, 3, 5)}$ does not carry any proper L-surgery structure positive or negative.*

Proof) Since $\Sigma(2, 3, 5) \# \overline{\Sigma(2, 3, 5)}$ does not have any tight contact structures positive or negative, it follows from Theorem 2.1 that $\Sigma(2, 3, 5) \# \overline{\Sigma(2, 3, 5)}$ does not have positive or negative L-surgery structures. $\square$

3 Ozsváth-Szabó's contact invariant

Here we review the Ozsváth-Szabó's contact invariant [7]. For a positive cooriented contact structure ξ over a closed oriented 3-manifold Y we will define the contact invariant $c(\xi)$ (originally defined in [7]). By E. Giroux's work [3] each isotopy class of contact structures exactly corresponds to an open book decomposition up to positive stabilization. An open book decomposition is a triple $D = (Y, K, \pi)$, where K is a fibered knot in Y and π is the fibration map $Y - K \rightarrow S^1$. The correspondence between an open book decomposition D and a contact structure is due to W. Thurston and H. Winkelnkemper [10]. We denote the correspondence as follows:

$$\{\text{open book decompositions}\} / \text{positive stabilization} \leftrightarrow \{\text{contact structures}\} / \text{isotopy}$$

$$D \rightarrow \xi_D.$$

Let (Y, η) be a contact structure and (Y, K, π) the open book decomposition of Y corresponding to η . We denote by the symbol $\mathfrak{t}(\eta)$ the spin^c structure associated with η . For $Y_0(K)$ there is the canonical contact structure ξ_0 satisfying $c_1(\xi_0)[\hat{F}] = 2g(\hat{F}) - 2$, where $\hat{F} \subset Y_0(K)$ is the capped surface of a fiber F of π . Then the natural spin^c cobordism $(-Y_0(K), \bar{\xi}_0) \rightarrow (-Y, \bar{\xi})$ induces $\widehat{HF}(-Y_0(K), \mathfrak{t}(\bar{\xi}_0)) \cong \mathbb{Z}_2 \oplus \mathbb{Z}_2 \rightarrow \widehat{HF}(-Y, \mathfrak{t}(\bar{\xi}))$. The invariant $c(\xi)$ is the image of the generator by the natural map $HF^+(-Y_0(K), \mathfrak{t}(\bar{\xi}_0)) \cong \mathbb{Z}_2 \rightarrow \widehat{HF}(-Y_0(K), \mathfrak{t}(\bar{\xi}_0))$. The main property used here is the following:

Theorem 3.1 ([7]) *If a positive contact structure (Y, ξ) is overtwisted, then $c(\xi) = 0$.*

From this theorem $c(\xi) \neq 0$ implies tightness of ξ .

4 Proof of Theorem 2.1

Proof) Let Y be an L-space homology sphere and $Y_p(K)$ an L-space. By the result in [8], we have $\widehat{HFK}(Y, K, g) \cong \mathbb{Z}_2$. Since $Y - K$ is irreducible, from the main theorem in [11], K is fibered. Hence we can build the contact structure over Y , which is associated with the open book decomposition $D = (Y, K, \pi)$ induced from this fibration. Consider the following surgery exact triangle.

$$\begin{array}{ccc}
 HF^+(-Y) & \xleftarrow{F_1} & HF^+(-Y_0(K), Q^{-1}[i]) \\
 & \searrow F_3 & \nearrow F_2 \\
 & HF^+(-Y_p(K), [i]) &
 \end{array} \tag{1}$$

Here $Q : \text{Spin}^c(-Y_0(K)) \cong \mathbb{Z} \rightarrow \text{Spin}^c(-Y_p(K))$ is a map between two sets of spin^c structures defined in [4]. Let $[i]$ denote the contact structure over $-Y_p(K)$ satisfying $Q(i) = [i]$. If all spin^c -structures in $Q^{-1}[i]$ are $c_1 \neq 0$, the map F_2 is the 0-map. Hence the map restricted of F_1 to the $\mathfrak{t}(\bar{\xi}_0)$ -component

$$HF^+(-Y_0(K), \mathfrak{t}(\bar{\xi}_0)) = HF^+(-Y_0(K), 1 - g) \cong \mathbb{Z}_2 \rightarrow HF^+(-Y)$$

is injective. Hence by definition of the contact invariant, $c(\xi_D)$ does not vanish. From the Theorem 3.1 ξ_D is a tight contact structure.

On the other hand in the case where $c_1(\mathfrak{t}(\bar{\xi}_0)) = 0$ the genus of K is one. Then for non-zero i , $HF^+(-Y_0(K), i) = 0$ and $HF_{\text{red}}(-Y_0(K), 0) = 0$. The knot Floer homology of K is $\widehat{HFK}(Y, K, i) \cong \mathbb{Z}_2$ for $i = 0, \pm 1$ and $\cong 0$ for $i \neq 0, \pm 1$. In this case the tau invariant $\tau(-K)$ is -1 by the same method as [8], hence the contact invariant $c(\xi_D)$ does not vanish. From Theorem 3.1 the contact structure ξ_D is tight. $\square$

We call a knot K in a homology sphere Y a *lens space Berge knot* if an integral Dehn surgery of K is a lens space and the dual knot K' of K is the union of two arcs each of which is embedded in the meridian disk of the genus one Heegaard decomposition (see Definition 1.7. in [8]).

The author has also verified that many Brieskorn spheres appear as the homology spheres Y yielding lens spaces. Ozsváth and Szabó have shown that any lens space Berge knot is fibered [8]. As a result many Brieskorn homology spheres carry proper L-surgery structure with contact structures associated with the lens space Berge knots. But I could not find proper L-surgery structures for Brieskorn homology spheres with the reversed orientation. Here we state a conjecture which generalizes Corollary 2.1.

Conjecture 4.1 *Let $\Sigma(p, q, r)$ be any Brieskorn homology sphere. Then $\Sigma(p, q, r)$ carries proper L-surgery structure but $\overline{\Sigma(p, q, r)}$ does not carry proper L-surgery structure.*

Acknowledgments

I thank Professor Ryosuke Yamamoto and T. Kadokami for their useful comments about contact structure and Dehn surgery respectively. It has inspired the author writing of the paper. I also thank Professor M. Ue for his giving me many suggestions in Doctor course seminar.

References

- [1] R. FINTUSHEL AND R. STERN, *Constructing lens spaces by surgery on knots*, Math. Z. Vol. 175, no. 1 February, (1980), 33-51
- [2] J. B. ETNYRE AND K. HONDA, *On the nonexistence of tight contact structures*, Annals of Mathematics **153**(2001) 749-766
- [3] E. GIROUX AND N. GOODMAN, *On the stable equivalence of open books in three-manifolds*, Geom. Topol. 10 (2006), 97-114
- [4] P. OZSVÁTH AND Z. SZABÓ, *Absolutely graded Floer homologies and intersection forms for four-manifolds with boundary*, Adv. Math. 173 (2003), no. 2, 179-261.
- [5] P. OZSVÁTH, Z. SZABÓ, *Holomorphic disks and topological invariants for closed three-manifolds*, Ann. of Math. (2) 159 (2004), no. 3, 1027-1158.
- [6] P. OZSVÁTH, Z. SZABÓ, *Holomorphic disks and three-manifold invariants: properties and applications*, Ann. of Math. (2) 159 (2004), no. 3, 1159-1245.
- [7] P. OZSVÁTH, Z. SZABÓ, *Heegaard Floer homologies and contact structures* Duke Math. J. **129**(2005) 39-61
- [8] P. OZSVÁTH AND Z. SZABÓ, *On knot Floer homology and lens space surgery*, Topology Volume 44, Issue 6 , November 2005, Pages 1281-1300
- [9] M. TANGE, *Lens spaces given from L-space homology 3-spheres* arXiv:0709.0141
- [10] W. THURSTON AND H. WINKELNKEMPER, *On the existence of contact forms*, Proc. A.M.S. **52** (1975), 345-347.
- [11] Y. NI, *Knot Floer homology detects fibered knots*, arXiv:math.GT/0607156

Motoo Tange
Research Institute for Mathematical Sciences,
Kyoto University,
Sakyo-ku Kyoto-shi Kyoto-fu 606-8502, Japan.
tange@kurims.kyoto-u.ac.jp